

SEARCHING FOR THE CAUSE OF DELIRIUM

Delirium Symptom list from CAM

- Sudden change in mental status
- Change in behaviour that fluctuates from normal to abnormal over a 24 hour period
- Difficulty in focusing attention
- Disorganized thinking and/or altered level of consciousness

Begin your assessment with the highest probable risk for your client's situation.

Drug Toxicity?

- a. On **more than six medications**, especially:
- anticonvulsants
 - histamine H₂ antagonist
 - insulin/hypoglycemic agent
 - antipsychotics
 - benzodiazepines
 - narcotics
 - barbiturates
 - thiazide diuretics
 - anticholinergics
 - antidepressants
 - cardiac glycosides
 - anesthetic
- b. **Receiving a medication for more than 5 years**
- c. **Age 75 or older**
- d. **Running drug levels beyond or at the high end of therapeutic range**
- Order drug chemistry and/or trial discontinuation of medicine.**

Changes in Chronic Illness?

Physical and psychosocial assessment reveals exacerbation* of previously diagnosed condition, such as:

- Diabetes mellitus
- COPD
- Cerebrovascular insufficiency
- Cancer
- Depression
- Substance misuse (e.g., alcohol, drugs, tobacco)
- Hypo/hypertension
- ASHD
- Pain
- Alzheimer disease/dementia
- Hypoxia

Request appropriate diagnostic tests

(* Exacerbation may be accompanied by increased levels of pain and/or decreased functional abilities)

New Disease Process?

- a) **Cardio and cerebrovascular conditions**
1. Silent MI
 2. TIA/CVA
 3. CHF
- or**
- b) **GI conditions**, GI bleed, if evidence of daily use of NSAIDs or steroids
- or**
- c) **Other medical conditions**
1. Hypo/hyperglycemia
 2. Hypo/hyperthyroidism
 3. Electrolyte imbalance
 4. Cancer
 5. Neurological conditions (e.g., normal pressure hydrocephalus)
 6. Pain
 7. Abuse or withdrawal from alcohol, drugs, tobacco
 8. Low B12
- Request appropriate diagnostic tests**
(e.g., PE, pulse oximetry, EKG, hemoglobin and hematocrit, chemistry screen, electrolytes, TSH, specific test for cancer detection, CAT)
- or**
- d) **Psychiatric conditions**, especially if evidence of family history
- Request psychiatric evaluation, dementia work up**

Adapted from:

- APA, 1987
- Inouye, SK, et al. (1990). Clarifying Confusion: The Confusion Assessment Method. A new method for detection of delirium. *Annals of Internal Medicine*, 113: 941-8.
- Inouye, SK. (2006). Delirium in older persons. *The New England Journal of Medicine*, 354(11), 1157-1165.
- Henry, M. (2002). Descending into delirium. *ANJ*, 102(3), p.49-56.
- Mentes, (1995) *Journal of Gerontological Nursing* in Henry, M. (2002, March). Descending into delirium. *ANJ*, 102(3), p.49-56.]

Infection?

- a. elevation in baseline temperature, even less than 37.56°C rectally
- b. history of lower respiratory infection or UTI more than twice per year
- c. history of any chronic infection
- d. recent episode of falling
- Request appropriate diagnostic tests.**
Most common: urinalysis, chest X-ray, sputum cultures

Elimination Problems?

- a. **Urinary problems**
- 1) history of incontinence, retention, or indwelling catheter
 - 2) signs or symptoms of dehydration, tenting, increased BUN
 - 3) decreased urinary output
 - 4) taking anticholinergic medication
 - 5) abdominal distention
- b. **Gastrointestinal problems**
- 1) immobility for more than 1 day in persons previously mobile
 - 2) abdominal distention
 - 3) decreased number of bowel movements or constipated stool
 - 4) decreased fluid intake – dehydration
 - 5) decreased food intake, especially bulk

Request in-out catheterization for postvoid residual and/or incontinence assessment, or both.
Accomplish digital rectal exam, request enema, initiate appropriate bowel regimen.

Sleep Disturbance?

- a) **Assess baseline normal sleep pattern**
- b) **Identify causes of sleep disturbance, e.g., Medications / pain / environment**

Post Operative?

- a) reaction to anesthetic
- b) analgesia
- c) opioids / anticholinergics
- Ensure "elder friendly"**

- a) Inactivity
- b) Restraint
- Mobilize early Manage pain**

Psychosocial? / Environmental?

- a) grief, losses (family members, significant life items)
- b) alteration in personal space
- c) recently admitted
- d) increase or decrease in sensory stimulation
- e) interpersonal difficulties

Initiate home assessment –

- a) ADLs and AIDLs
- b) Safety
- c) User-friendly environment
- labels, pictures
 - put orienting items in room
- d) Supports: social, family; counseling
- e) Encourage family involvement